

TANGENTIAL CENTER PROBLEM IN DIMENSION ZERO

A. ÁLVAREZ, J.L. BRAVO, P. MARDEŠIĆ

ABSTRACT. This paper is dedicated to the solution of the tangential center problem on 0-cycles:

Given a polynomial $f \in \mathbb{C}[z]$ of degree m , denote $z_1(t), \dots, z_m(t)$ all algebraic functions defined by $f(z_k(t)) = t$. Given integers $n_1, \dots, n_m$ such that $n_1 + \dots + n_m = 0$, find all polynomials $g \in \mathbb{C}[z]$ such that $n_1 g(z_1(t)) + \dots + n_m g(z_m(t)) \equiv 0$.

The classical Center-Focus Problem, or rather its tangential version in important non-trivial planar systems lead to the above problem.

We prove the uniqueness of decompositions $f = f_1 \circ \dots \circ f_d$, such that every f_k is 2-transitive, monomial or a Chebyshev polynomial under the assumptions that in the above composition there is no merging of critical values.

Under these assumptions, we give a complete (recursive) solution of the tangential center problem on 0-cycles. The recursive solution is obtained through three mechanisms: composition, primality and vanishing of the Newton-Girard component on projected cycles.

The first mechanism was studied by Pakovich and Muzychuk and the authors among others in previous publications. The two later mechanisms are new.

1. INTRODUCTION

A classical problem in planar polynomial vector fields starting from Poincaré is the *center-focus problem*. The problem asks for the determination of mechanisms leading to the creation of a center (a singularity surrounded by a continuous family of closed orbits) rather than a focus (singularity attracting or repelling all nearby trajectories). The problem has not yet been solved in a satisfactory way except for quadratic vector fields [12], [16].

The *center-focus problem* has its *infinitesimal version*: Starting from a vector field X_0 having a center, find all perturbations X_λ preserving the center. More generally, given a vector field X_0 having a continuous family $\gamma_0(t)$ of closed orbits, determine the deformations X_λ of X_0 preserving these closed orbits. That is, we ask the deformed family $\gamma_\lambda(t)$ of orbits of X_λ to be *closed*, too.

Taking a parametrized transversal T to the family of closed curves $\gamma(t)$, one defines the *displacement map* Δ as the first return map minus identity along trajectories of X_λ . Then, a continuous family of closed curves is preserved, if the displacement map of the deformation along the chosen family of closed curves is identically zero.

The most popular family of polynomial systems having a continuous family of closed orbits is the family of Hamiltonian vector fields $X_F = -\frac{\partial F}{\partial y} \frac{\partial}{\partial x} + \frac{\partial F}{\partial x} \frac{\partial}{\partial y}$. The trajectories of X_F lie in the level curves of the Hamiltonian. Consider their deformations $X_F + \epsilon Y$. Taking a transversal T to a family $\gamma(t)$ of closed curves of the Hamiltonian vector field X_F parametrized by F , the displacement function of the above deformation is of the form

$$\Delta(t, \epsilon) = -\epsilon \int_{\gamma(t)} \omega_Y + o(\epsilon),$$

where ω_Y is the dual form to the vector field Y and $o(\epsilon)$ is a function depending on t , but tending to zero faster than ϵ , for $\epsilon \rightarrow 0$. The function $t \mapsto I(t) = \int_{\gamma(t)} \omega_Y$, $\gamma(t) \subset F^{-1}(t)$, is an *abelian integral*.

A necessary condition for having a solution of the infinitesimal center problem is the vanishing of its first-order term given by the abelian integral $I(t)$.

This motivates the *tangential (or first-order) center problem*. We formulate it in its complex form.

Problem 1.1. Tangential center problem. *Given a polynomial $F \in \mathbb{C}[x, y]$ and a continuous family $\gamma(t)$ of cycles belonging to the first homology group $H_1(F^{-1}(t))$, find all polynomial forms ω such that the abelian integral*

$$(1.1) \quad I(t) = \int_{\gamma(t)} \omega$$

vanishes identically.

The problem was solved by Ilyashenko [15] under a genericity assumption on F . He proves that under a genericity condition on F , for any family of cycles $\gamma(t) \in H_1(F^{-1}(t))$, the abelian integral (5.14) vanishes identically if and only if the form ω is relatively exact, i.e., of the form $\omega = GdF + dR$, for some $G, R \in \mathbb{C}[x, y]$.

Without the genericity hypothesis on F , the claim is false and the tangential center problem is open. One important non-generic case is the *hyper-elliptic case* when $F(x, y) = y^2 + f(x)$, $f \in \mathbb{C}[x]$. For hyper-elliptic F , the problem was solved by Christopher and the third author [10] under the hypothesis that the family of cycles $\gamma(t)$ is a family of vanishing cycles. The tangential-center problem is open even in the hyper-elliptic case for general family $\gamma(t)$ of cycles.

Let us note a related problem studied by Gavrilov [13] and Bonnet and Dimca [3].

Problem 1.2. Integrability problem. *Given a polynomial $F \in \mathbb{C}[x, y]$, find all polynomial forms ω such that the abelian integral $I(t) = \int_{\gamma(t)} \omega$ vanishes identically along any cycle $\gamma(t) \in H_1(F^{-1}(t))$.*

Note the difference: in the Integrability problem one asks for the vanishing of abelian integrals along *all* cycles and not just one family of cycles as in the Tangential center problem. Under generic hypothesis on F , Ilyashenko [15] proves that by monodromy one family of cycles generates all cycles. Hence the vanishing of abelian integrals along one family of cycles implies the vanishing along all cycles and the two problems coincide.

In general it is false. It is easy to see that there are solutions of the tangential center problem due to the presence of a symmetry on a family of cycles. This solution will not necessarily be a solution for another family of cycles not respecting the symmetry. The form is hence not relatively exact as Abelian integrals of relatively exact forms vanish along any cycle of $\gamma(t)$.

This paper is dedicated to the study of the 0-dimensional version of the tangential center problem:

Problem 1.3. 0-dimensional tangential center problem. *Given a polynomial $f \in \mathbb{C}[x]$ and a family of 0-cycles $C(t)$ of f , determine all functions $g \in \mathbb{C}[x]$ such that $\int_{C(t)} g$ vanishes identically.*

Here, integration is just calculation of the value of a function at some points $z_i(t)$ determining the cycle $C(t) = \sum n_i z_i(t)$ (for more details see the next section).

Note that in this case 0-dimensional abelian integrals are in fact algebraic functions, so the problem is certainly easier than the initial problem.

Nevertheless, in [2], we showed that the tangential center problem for the hyper-elliptic case is directly related to the above 0-dimensional tangential center problem for hyper-elliptic abelian integrals, (see also Gavrilov and Pakovich [14]).

In [2] we introduced the classes of *balanced and unbalanced cycles*. We solved the 0-dimensional tangential center problem by induction under the hypothesis that the initial cycle is unbalanced and that in the inductive process one encounters only unbalanced cycles (see also [14]). We called such a cycle *totally unbalanced*. The main result from [2] can be resumed by saying that under the hypothesis that the cycle $C(t)$ is totally unbalanced, the only mechanism producing tangential centers is a composition (i.e. symmetry) mechanism or a sum of composition mechanisms. Our proof was based on results of Pakovich-Muzychuk relative to the composition conjecture in the moment problem for the Abel equation. In particular we used Pakovich-Muzychuk's characterization of invariant irreducible spaces.

In this paper we deal with the remaining case of balanced cycles or cycles leading to balanced cycles in the induction process. We solve by induction the 0-dimensional tangential center problem for arbitrary cycles, but under a hypothesis on the polynomial f . We explain this hypothesis. It follows from the Burnside-Schur theorem that primitive (i.e., undecomposable) composition factors of a polynomial f are of one of the following three types: 2-transitive, linearly equivalent to a monomial z^k or a Chebyshev polynomial T_k , with k prime. Note that a composition of monomials is a monomial and similarly a composition of Chebyshev polynomials is a Chebyshev polynomial.

Given a polynomial f let $f = f_0 \circ \dots \circ f_d$ be a decomposition of the polynomial f in its composition factors, which are 2-transitive, Chebyshev or monomial *not necessarily of prime degree*.

We completely solve the tangential center problem in dimension zero when for every $k = 1 \dots, d$ the critical values of $f_0 \circ \dots \circ f_{k-1}$ and f_k do not merge in the composition $f_0 \circ \dots \circ f_k$ (see Definition 2.5). We prove that under the above hypothesis, the decomposition of f as $f = f_0 \circ \dots \circ f_d$ is unique. The hypothesis of non-merging of critical values allow us to decompose the monodromy group of $f_0 \circ \dots \circ f_k$ in a semidirect product of the monodromy group of $f_0 \circ \dots \circ f_{k-1}$ and the monodromy group of f_k . Under these hypotheses, we solve recursively the 0-dimensional tangential center problem. Using new methods, we show that in addition to the composition mechanisms, two new mechanisms leading to tangential centers exist. They both appear for balanced cycles on the level of the basis of recursion and in the recursion step.

One mechanism is related to primality of monomial or Chebyshev polynomial factors with the characteristic polynomial $P_C(z) = \sum_{i=1}^m n_i z^{i-1}$ associated to the cycle $C(t) = \sum_{i=1}^m n_i z_i(t)$. It is described by (1) of Theorem A (basis) and Theorem C (recursive step). The other mechanism is related to the vanishing of the Newton-Girard component on projected cycles in the case of 2-transitive factors. It is described by (2) of Theorem A (basis) and Theorem B (recursive step).

The motivation of the tangential center problem for the Abel equation

$$(1.2) \quad x' = p(t)x^2 + \lambda q(t)x^3,$$

proposed by Briskin, Françoise and Yomdin in a series of papers [5]-[7], was to obtain general mechanisms for the existence of centers. For p, q polynomials, the tangential center problem has been totally solved by Pakovich and Muzychuk [19] proving that there is a tangential center if and only if $\int_0^t q(s) ds$ can be written as a sum of polynomials having a common factor with $\int_0^t p(s) ds$. Moreover, Briskin,

Roytvarf and Yomdin [8] have proved that composition (just one summand) generates all centers of (1.2) at infinity except for a finite number of exceptional cases. Nevertheless, there are centers and tangential centers for which composition is not the generating mechanism, like (1.2) when p, q are not polynomials (see [1, 9]) or, in a more general context, (hyperelliptic) planar systems (see Example 9.2 of [2]). The new mechanisms obtained here explain the hyperelliptic planar system case and we hope they will shed some light on the general moment problem.

2. MAIN RESULTS

Let $f \in \mathbb{C}[z]$ be a polynomial of degree m . Points $z \in \mathbb{C}$ such that $f'(z) = 0$ are critical points of f . Corresponding values $t = f(z)$ are critical values and non-critical values are called regular values. Let Σ denote the set of critical values of f . Then $f : f^{-1}(\mathbb{C} \setminus \Sigma) \rightarrow \mathbb{C} \setminus \Sigma$ is a fibration with 0-dimensional fiber. For any regular value of t the fiber $f^{-1}(t)$ consists of m distinct points. These points can be continuously transported along any path in $\mathbb{C} \setminus \Sigma$. Replacing in the fibration $f : f^{-1}(\mathbb{C} \setminus \Sigma) \rightarrow \mathbb{C} \setminus \Sigma$ the fibers $f^{-1}(t)$ by their 0-th homology groups $H_0(f^{-1}(t))$ or their reduced 0-th homology groups $\tilde{H}_0(f^{-1}(t))$, one obtains the homology or reduced homology fibration.

Let t_0 be a regular value of f and $\pi_1(\mathbb{C} \setminus \Sigma, t_0)$ the first homotopy group of the base with base point t_0 . Then the continuous transport gives a morphism $\pi_1(\mathbb{C} \setminus \Sigma, t_0) \rightarrow \text{Aut}(H_0(f^{-1}(t_0)))$ from the first homotopy group of the base to the group of automorphisms of the homology fiber. Its image is called the monodromy group G_f of f . This group is isomorphic to the Galois group of $f(z) - t$ seen as a polynomial over $\mathbb{C}(t)$ (see [10, Th. 3.3], for example). Any section C of the homology fibration is called a 0-chain. A section of the reduced homology fibration is called a 0-cycle.

Choosing an order $\{z_1(t), \dots, z_m(t)\}$ among the m -preimages of t by f , a 0-chain of f is represented by

$$(2.3) \quad C(t) = \sum_{i=1}^m n_i z_i(t), \quad n_i \in \mathbb{Z}$$

and 0-cycles moreover satisfy

$$(2.4) \quad \sum_{i=1}^m n_i = 0.$$

Note that $z_i(t)$, as well as $C(t)$ are multivalued functions on $\mathbb{C} \setminus \Sigma$. We also consider chains with coefficients in $\mathbb{C}$ when necessary. Then we will specify the coefficient ring.

Let C be a chain of f and g a polynomial. The 0-dimensional Abelian integral of g along C is defined by

$$\int_{C(t)} g := \sum_{i=1}^m n_i g(z_i(t)).$$

This paper is dedicated to the solution of Problem 1.3 above.

In [2] we introduced the notions of balanced and unbalanced cycles. We proved that if the cycle is unbalanced, then the tangential center problem is equivalent to solving some induced tangential center problems with f replaced by some composition factors of f , hence polynomials of smaller complexity and smaller degree. Let us precise the notions. In order to be able to perform the induction we need to extend the notions from cycles to chains.

Let $\Gamma_m(f) \subset G_f$ denote the conjugacy class of τ_∞ , where $\tau_\infty \in G_f$ is associated to a path winding once counter-clockwise around infinity. That is, $\Gamma_m(f)$ is the

set of all $\sigma \circ \tau_\infty \circ \sigma^{-1}$, for any $\sigma \in G_f$. We label the roots so that τ_∞ , which is a permutation cycle of order m , shifts the indices of the roots by one, i.e., $\tau_\infty = (1, 2, \dots, m)$.

Definition 2.1. We say that a chain $C(t)$ of f is balanced if

$$\sum_{i=1}^m n_{p_i} \epsilon_m^i = 0, \quad \text{for every } \tau = (p_1, p_2, \dots, p_m) \in \Gamma_m(f),$$

where ϵ_m is any primitive m -th root of unity. If $C(t)$ is not balanced, we say that $C(t)$ is unbalanced.

We call the polynomial

$$(2.5) \quad P_C(z) = \sum_{n=1}^m n_j z^{j-1}$$

characteristic polynomial of the chain $C(t)$.

Note that if $C(t)$ is balanced, then $P_C(\epsilon_m) = 0$. The converse does not hold in general since the characteristic polynomial depends on the election and of the representation of the cycle of infinity, so indeed we should define the characteristic ideal as in [2], but in the special cases we use the characteristic polynomial, $C(t)$ is balanced if and only if $P_C(\epsilon_m) = 0$.

The notion of balanced chain is independent on the way how permutation cycles are written. Let $\tau \in \Gamma_m(f)$ be of the form $\tau = \sigma \circ \tau_\infty \circ \sigma^{-1}$, $\sigma \in G_f$. Put $(p_1, p_2, \dots, p_m) = (\sigma(1), \sigma(2), \dots, \sigma(m))$ and

$$(n_{p_1}, n_{p_2}, \dots, n_{p_m}) = (n_{\sigma(1)}, n_{\sigma(2)}, \dots, n_{\sigma(m)}) = \sigma(n_1, n_2, \dots, n_m).$$

Then, we can rephrase the definition of a chain $C(t)$ being *balanced* as

$$\sum_{i=1}^m n_{\sigma(i)} \epsilon_m^i = 0, \quad \text{for every } \sigma \in G_f.$$

Assume that $f = \tilde{f} \circ h$, for $\tilde{f}, h \in \mathbb{C}[z]$, $\deg(h) = d$. Consider the imprimitivity system $\mathcal{B} = \{B_k\}_{k=1, \dots, m/d}$, where $B_k = \{i \in \{1, 2, \dots, m\} | h(z_i(t)) = h(z_k(t))\}$, associated to h .

Definition 2.2. The cycle $h(C(t))$ of $\tilde{f}$ called the projection of $C(t)$ is defined by

$$(2.6) \quad h(C(t)) = \sum_{h(z_i(t))} \left(\sum_{h(z_j)=h(z_i)} n_j \right) h(z_i(t)) = \sum_{k=1}^{m/d} \left(\sum_{i \in B_k} n_i \right) w_k(t).$$

Here $w_1(t), \dots, w_{m/d}(t)$ are all the different roots $h(z_i(t))$ of $\tilde{f}(z) = t$.

With the above notation, the main result of [2]: Theorem 2.2 (ii) can be stated as follows.

Theorem 2.3 ([2]). Let $f \in \mathbb{C}[z]$, and let $C(t)$ be an unbalanced cycle. Then,

$$\int_{C(t)} g \equiv 0, \quad \text{for } g \in \mathbb{C}[z]$$

if and only if there exist $f_i, g_i, h_i \in \mathbb{C}[z]$, $1 \leq i \leq d$ such that $\deg(h_i) > 1$, $f = f_i \circ h_i$, $1 \leq i \leq d$, $g = \sum_{i=1}^d g_i \circ h_i$ and $\int_{h_i(C(t))} g_i \equiv 0$.

Remark 2.4. If the cycle $C(t)$ is unbalanced, then the tangential center problem reduces to solving it for the projected cycles. If some of these projected cycles are unbalanced, Theorem 2.3 applies again. In particular, if all projected cycles (in every successive step) are unbalanced, we call the initial cycle totally unbalanced and the problem is completely solved by induction using Theorem 2.3.

In this paper we study the remaining case, when $C(t)$ is balanced or in the progress of projecting we arrive at some balanced cycle. It appears that the decompositions of the polynomial f play a central role in the problem.

The action of the monodromy group G_f of f on the set of solutions $z_i(t)$ of the equation $f(z) = t$ is closely related to the decomposability of f . When a group G acts on a finite set X , it is said that the action is *imprimitive* if X can be non-trivially decomposed into subsets of the same cardinality B_i such that every $g \in G$ sends each B_i into B_j for some j . Otherwise, the action is called *primitive*. The action of G is said to be *2-transitive* if given any two pairs of elements of X , (i, j) and (k, l) , there is an element $g \in G$ such that $\sigma(i) = k$ and $\sigma(j) = l$. It is easy to prove that in these cases X cannot be divided into disjoint subsets such that G acts on them, which is one of the trivial cases of a primitive action. These disjoint sets B_i are called *blocks* and they form a partition of X called an *imprimitivity system*. Observe that by [2, Prop. 4.1] this definition is consistent with the previous one.

Precisely, the action of G_f on $\{z_i(t)\}$ is imprimitive if and only if f is decomposable (see [10, Prop. 3.6]). Burnside-Schur Theorem (see e.g. [10, Th. 3.8]) classifies primitive polynomials (i.e., a polynomial that cannot be written as a composition of two polynomials of degree greater than one):

Burnside-Schur Theorem ([10, Theorem 3.8]) *Let f be a primitive polynomial and G_f its monodromy group. Then, one of the following holds:*

- (1) *The action of G_f on the $z_i(t)$ is 2-transitive. We call such a polynomial 2-transitive.*
- (2) *f is (linearly) equivalent to a Chebyshev polynomial T_p where p is prime.*
- (3) *f is (linearly) equivalent to z^p where p is prime.*

We give a solution of the tangential center problem by induction on the number of composition factors f_i under some additional hypotheses. We first solve the problem for basic composition factors: 2-transitive, monomial or Chebyshev of *not necessarily prime degree* (basis of induction). Monomial and Chebyshev cases are similar. They are hence treated together.

We introduce first some notations. Given a polynomial f of degree m and $z_i(t)$, $i = 1, \dots, m$ the roots of $f(z) = t$, denote s_k , $k = 0, \dots, m-1$ the sums of k -th powers of roots of $f - t$:

$$s_k = \sum_{i=1}^m z_i^k(t).$$

Note that the Newton-Girard formulae express $s_0, \dots, s_{m-1}$ in function of the coefficients of f . In particular they are independent of t . Denote $s(f) = (s_0, \dots, s_{m-1}) \in \mathbb{C}^m$ the Newton-Girard vector of f .

When convenient, using the division algorithm, we will write $g(z)$ as $g(z) = \sum_{k=0}^{m-1} g_k(f)z^k$. That is we express $g(z)$ as a polynomial in z of degree $\deg g < m = \deg f$, with coefficients in $C[f]$. Substituting $f = t$ for integration each coefficient g_k becomes a polynomial $g_k(t)$ in t .

We denote $g(t) = (g_0(t), \dots, g_{m-1}(t))$. Let $\langle, \rangle$ denote the scalar product in $\mathbb{C}^m$: $\langle g(t), s(f) \rangle = \sum_{i=0}^{m-1} g_i(t)s_i$.

We will also use cyclotomic polynomials. We denote $\Phi_j(z)$ the j -th cyclotomic polynomial given by

$$\Phi_j(z) = \prod_{\substack{1 \leq j \leq m, \\ \gcd(j, m)=1}} (z - e^{\frac{2\pi i}{m}j}).$$

Theorem A. *Assume that $C(t) = \sum n_j z_j(t)$ is a balanced chain of a polynomial f .*

- (1) If the action of G_f is 2-transitive, then in the orbit of $C(t)$ there exists a cycle $D(t) = \sum n_j z_j(t)$ such that $n_j = n$ for every $1 \leq j \leq m$. Moreover, for every $g(z) = \sum_{k=0}^{m-1} g_k(f) z^k$, $\int_{C(t)} g \equiv 0$ if and only if $g(t) \in s(f)^\perp$.
- (2) If $f(z) = z^m$ (resp. $f(z) = T_m(z)$), then $\int_{C(t)} g \equiv 0$ if and only if $g(z) = \sum_{j=0}^{m-1} g_j(h_j(z))$, where $g_j \in \mathbb{C}[z]$, $h_j(z) = z^j$ (resp. $h_j(z) = T_j(z)$), for any j is such that $\Phi_{m/k}(z) | P_C(z)$, where $k = \text{g.c.d.}(m, j)$, $P_C(z) = \sum_{i=1}^m n_i z^{i-1}$ is the characteristic polynomial of the chain C and $\Phi_j(z)$ is the j -th cyclotomic polynomial.

Condition (2) means that the only terms h_j present in the solution $g = \sum_{j=0}^{m-1} g_j h_j(z)$ of the tangential center problem for f monomial or Chebyshev polynomial of degree m are $j|m$ such that $P_C(\epsilon_m^j) = 0$, where ϵ_m is an m -th root of unity.

Next we show how to reduce the tangential center problem on a cycle $C(t)$ of $f = \tilde{f} \circ h$, where h is 2-transitive or linearly equivalent to a monomial or a Chebyshev polynomial of not necessarily prime degree. In the step of induction, we reduce the original tangential center problem of a cycle of f to the tangential center problem of the projected cycle $h(C(t))$, which is a cycle of the function $\tilde{f}$, simpler than the original function f . We do the induction step under the generic hypothesis that the critical values of $\tilde{f}$ and h do not merge:

Definition 2.5. Let $\tilde{f}$ and h be two nonlinear polynomials and let $f = \tilde{f} \circ h$. We say that the critical values of $\tilde{f}$ and h do not merge if

- (1) $\{f(z) : \tilde{f}'(h(z)) = 0\} \cap \{f(z) : h'(z) = 0\} = \emptyset$,
- (2) $\tilde{f}$ is injective on the set of critical values of h .

Condition (1) of Definition 2.5 assures that the monodromy group G_f is a semidirect product of the subgroups N_h and $G_{\tilde{f}}$, $G_f = N_h \rtimes G_{\tilde{f}}$, where N_h is the normal closure of the monodromy group $G_h \subset G_f$ with the natural injection (which in general is not a group morphism). Condition (2) of Definition 2.5 assures that there exist group morphisms $\phi : G_h \rightarrow G_f$.

We divide the induction step in two cases, according to the nature of h . We study first the case when h is 2-transitive. Next the case when h is a monomial or a Chebyshev polynomial.

Theorem B. Let $f(z)$ be a polynomial such that $f = \tilde{f} \circ h$ with $\tilde{f}, h \in \mathbb{C}[z]$, $1 < \deg(h) = d < m = \deg f$. Let $C(t) \in \tilde{H}_0(f^{-1}(t))$ be a balanced cycle of f . Assume that the critical values of $\tilde{f}$ and h do not merge and that let h be 2-transitive.

Let $\tilde{z}_k(w)$, $k = 1, \dots, d$, denote the zeros of $h(z) - w$ and let $s_i = \sum_{k=1}^d \tilde{z}_k^i(w)$, $i = 0, \dots, d-1$ be the sum of powers of roots of $h(z) - w$.

Then $g(z) = \sum_{i=0}^{d-1} z^i g_i(h(z))$, verifies

$$(2.7) \quad \int_{C(t)} g \equiv 0$$

if and only if

$$(2.8) \quad \int_{h(C(t))} \tilde{g} \equiv 0,$$

and $g_i \in \mathbb{C}[w]$, $i = 0, \dots, m-1$, are solutions of the linear system

$$(2.9) \quad \sum_{i=0}^{d-1} s_i g_i(w) = \tilde{g}(w).$$

Remark 2.6. Thus, the solution g and the inductive solution $\tilde{g}(w)$ are related by the linear system of equations $\tilde{g}(w) = \sum_{i=0}^{d-1} s_i g_i(w)$. Here s_i , $i = 0, \dots, d-1$ are expressed through the coefficients of h by the Newton-Girard formulae, for $h(z) - w$ and are independent of w . We call $s(h) = (s_0, \dots, s_{d-1}) \in \mathbb{C}^d$ the Newton-Girard vector of h . Let $\pi_h(g) = \frac{\langle g(w), s(h) \rangle}{|s(h)|}$ be the component of the vector $g(w) = (g_0(w), \dots, g_{d-1}(w)) \in \mathbb{C}[w]^d$ representing g in the direction of the Newton-Girard vector $s(h)$. Theorem B can be resumed by saying that under above conditions (6.18) is equivalent to the vanishing of $\int_{h(C(t))} \pi_h(g)$, where $\pi_h(g)$ is the component of the vector representing g in the direction of the Newton-Girard vector of h . See Example 2.10.

Assume that $f = \tilde{f} \circ h$, for $\tilde{f}, h \in \mathbb{C}[z]$, $\deg(h) = d$. In order to formulate the step of recursion in the case of monomial or Chebyshev h , (Theorem C), in addition to the projection of a cycle $C(t)$ by h we need the notion of h -invariant parts of $C(t)$:

Definition 2.7. For each cycle $C(t)$ of f and the decomposition $f = \tilde{f} \circ h$ of f , consider the imprimitivity system $\mathcal{B} = \{B_1, \dots, B_{m/d}\}$ associated to h . We define the h -invariant parts of $C(t)$ as the chains $C_k(t) = \sum_{i \in B_k} n_i z_i(t)$ of f , or as chains $\tilde{C}_k(w) = \sum_{i \in B_k} n_i z_i(w)$, $h(z_i) = w$, of h .

The h -invariant part $C_k(t)$ corresponds to the k -th line in Figure 4. Relations $C(t) = \sum_{k=1}^{m/d} C_k(t)$ and $\tilde{C}_k(w) = C_k(\tilde{f}(w))$ hold. Note that even if $C(t)$ is a cycle, the h -invariant parts $C_k(t)$ are only chains in general.

Theorem C. Let $f(z)$ be a polynomial such that $f = \tilde{f} \circ h$ with $\tilde{f}, h \in \mathbb{C}[z]$, $1 < \deg(h) = d < m = \deg f$. Let $C(t) \in \tilde{H}_0(f^{-1}(t))$ be a balanced cycle of f . Assume that the critical values of $\tilde{f}$ and h do not merge and let h be a monomial or a Chebyshev polynomial. Let $\tilde{z}_k(w)$, $k = 1, \dots, d$, denote the zeros of $h(z) - w$. Then the h -invariant parts $\tilde{C}_k(w)$, $k = 1, \dots, m/d$, of $C(t)$ are balanced.

Moreover,

$$\int_{C(t)} g \equiv 0$$

if and only if g is of the form

$$g(z) = \tilde{g}(h(z))/(d-1) + u(z),$$

where $\tilde{g}(w)$ is a polynomial such that

$$(2.10) \quad \int_{h(C(t))} \tilde{g} \equiv 0,$$

and

$$(2.11) \quad \int_{\tilde{C}_k(w)} u \equiv 0, \quad 1 \leq k \leq m/d.$$

Note that as $\tilde{C}_k(w)$ are balanced chains of h , (2.11) is solved in Theorem A and the solution of (6.16) is given by the induction hypothesis.

We prove Theorems B and C in Section 6.

If $\tilde{f}$ and h merge, then Theorems B and C also gives tangential centers. However, there can be other solutions not covered by them. Indeed, by perturbing $f = \tilde{f} \circ h$ we can assure that $\tilde{f}$ and h do not merge and Theorems B and C gives solutions of the deformed system, which in the limit give solutions of the original system.

It follows from the Burnside-Schur Theorem that any polynomial f can be decomposed as $f = f_0 \circ \dots \circ f_d$, where each factor f_k is either 2-transitive, linearly equivalent to a monomial or linearly equivalent to a Chebyshev polynomial. Note

that a composition of monomials is a monomial and a composition of Chebyshev polynomials is a Chebyshev polynomial. We do not assume that the degree of these polynomials f_k linearly equivalent to a monomial or to a Chebyshev polynomial is prime. Putting together Theorem 2.3 and Theorems A, B and C, we solve inductively the tangential center problem under the following hypothesis on f :

Hypothesis 2.8. *Let $f \in \mathbb{C}[z]$ have a decomposition $f = f_0 \circ \dots \circ f_d$, with f_k 2-transitive, linearly equivalent to a monomial or to a Chebyshev polynomial. We assume that the critical values of $f_0 \circ \dots \circ f_{k-1}$ and f_k do not merge for any $1 \leq k \leq d$.*

We show that under Hypothesis 2.8, there exists a unique decomposition $f = f_0 \circ \dots \circ f_d$, with f_k 2-transitive, linearly equivalent to a monomial or to a Chebyshev polynomial, satisfying Hypothesis 2.8. More precisely:

Proposition 2.9. *Assume that $f = f_0 \circ \dots \circ f_d$ for some polynomials $f_0, \dots, f_d$ such that every f_k is either 2-transitive, linearly equivalent to a monomial or linearly equivalent to a Chebyshev polynomial, and that Hypothesis 2.8 holds.*

If $f = \tilde{f}_0 \circ \dots \circ \tilde{f}_{\tilde{d}}$ is another decomposition of f for some polynomials $\tilde{f}_0, \dots, \tilde{f}_{\tilde{d}}$ such that every $\tilde{f}_k$ is either 2-transitive, linearly equivalent to a monomial or linearly equivalent to a Chebyshev polynomial, then there exist $j_{k,1}, \dots, j_{k,d(k)}$, $j_{k,1} = j_{k-1,d(k-1)} + 1$ ($k > 0$), such that (up to linear transformations)

$$f_k = \tilde{f}_{k,1} \circ \dots \circ \tilde{f}_{k,d(k)}, \quad \text{for every } k = 0, 1, \dots, d.$$

Moreover, if the critical values of $\tilde{f}_0 \circ \dots \circ \tilde{f}_{\tilde{d}-1}$ and $\tilde{f}_{\tilde{d}}$ do not merge for any $1 \leq k \leq \tilde{d}$, then $d = \tilde{d}$, and (up to linear transformations)

$$f_k = \tilde{f}_k, \quad \text{for every } k = 0, 1, \dots, d.$$

To conclude this section, we give two examples of the application of Theorems A, B and C.

Example 2.10. *First, let us consider a polynomial composition of two 2-transitive polynomials $f = \tilde{f} \circ h$, with $\tilde{f}(z) = z^3 - z^2 + z$ and $h(z) = z^3 + 2z^2 - 1$, and the cycle*

$$C(t) = z_1(t) - z_2(t) + z_4(t) - z_5(t) + z_7(t) - z_8(t),$$

where we are assuming that $(1, 2, 3, 4, 5, 6, 7, 8, 9)$ is the permutation associated to a loop around infinity. It can be checked that Hypothesis 2.8 holds. Moreover, $C(t)$ is balanced.




FIGURE 1. Cycles $C(t)$ and $h(C(t))$ in Example 2.10

Write $g(z) = g_0(h(z)) + zg_1(h(z)) + z^2g_2(h(z))$, $g_i \in \mathbb{C}[w]$. By Theorem B, g satisfies $\int_{C(t)} g \equiv 0$ if and only if

$$\int_{h(C(t))} \tilde{g} \equiv 0,$$

where $h(C(t)) = w_1(t) - w_2(t)$ is unbalanced, and $\tilde{g}(w) = 3g_0(w) - 2g_1(w) + 4g_2(w)$.

Now, the solutions of $\int_{h(C(t))} \tilde{g} \equiv 0$ are given by Theorem 2.3. As $\tilde{f}$ is primitive, the solutions are of the form $\tilde{g}(w) = k_0(\tilde{f}(w))$, for any polynomial $k_0 \in \mathbb{C}[w]$.

Then $\int_{C(t)} g \equiv 0$ if and only if $k_0(\tilde{f}(w)) = 3g_0(w) - 2g_1(w) + 4g_2(w)$. For instance, if $k_0(w) = 1$, and g_0, g_1, g_2 are quadratic polynomials: $g_i(z) = a_{i2}z^2 + a_{i1}z + a_{i0}$. Then the space of solutions is six-dimensional given by

$$\begin{aligned} g(z) = & 1 + a_{10} \left(\frac{2}{3} + z \right) + a_{11} \frac{1+z}{3} (2+3z)(-1+z+z^2) \\ & + a_{12} \frac{2+3z}{3} (-1+2z^2+z^3)^2 \\ & + a_{20} \left(-\frac{4}{3} + z^2 \right) + a_{21} \frac{1+z}{3} (-1+z+z^2)(-4+3z^2) \\ & + a_{22} \frac{-4+3z^2}{3} (-1+2z^2+z^3)^2, \end{aligned}$$

for any $a_{10}, a_{11}, a_{12}, a_{20}, a_{21}, a_{22} \in \mathbb{C}$.

Example 2.11. Let us consider a polynomial composition of a 2-transitive polynomial and a monomial, $f = \tilde{f} \circ h$, with $\tilde{f}(w) = w^3 - w^2 + w$ and $h(z) = z^6$, and the cycle

$$C(t) = z_1(t) - z_2(t) + z_7(t) - z_8(t) + z_{13}(t) - z_{14}(t),$$

where we are assuming that $(1, 2, \dots, 17, 18)$ is the permutation associated to a loop around infinity. It can be checked that Hypothesis 2.8 holds. Moreover, $C(t)$ is balanced.

It is easy to see that $h(C(t)) = 3w_1(t) - 3w_2(t)$ is an unbalanced cycle of $\tilde{f}$. Then, the solutions of

$$\int_{h(C(t))} \tilde{g} \equiv 0$$

are given by Theorem 2.3. More precisely, as $\tilde{f}$ is primitive, it follows that $\tilde{g}(w)$ is a function of $\tilde{f}$. Denote $\tilde{g}(w) = 5g_0(\tilde{f}(w))$. On the other hand, the h -invariant parts of the cycle $C(t)$ are given by $\tilde{C}_1(t) = w_1(t) + w_3(t) + w_5(t)$, $\tilde{C}_2(t) = -w_1(t) - w_3(t) - w_5(t)$ and $\tilde{C}_3(t) = 0$. Hence $P_{\tilde{C}_1}(w) = 1 + w^2 + w^4$, $P_{\tilde{C}_2} = -P_{\tilde{C}_1}$ and $P_{\tilde{C}_3} = 0$.

if $u = \sum u_k z^k$, by Theorem A (2), the solutions of

$$\int_{\tilde{C}_k(t)} u \equiv 0, \quad \text{for every } k,$$

are given by $u_{3k} = 0$ for every $k = 0, 1, 2, \dots$

Finally, by Theorem C, the solutions of

$$\int_{C(t)} g \equiv 0$$

are

$$g(z) = g_0(f(z)) + u(z),$$

for any polynomials $g_0, u = \sum u_k z^k$ such that $u_{3k} = 0$ for every $k = 0, 1, 2, \dots$

3. STRUCTURE OF THE SPACE OF SOLUTIONS

In this section we give the general structure of the space of solutions. It will be used in the next section for solving the tangential center problem in two basic cases: the monomial and the Chebyshev case.

Choosing a basis $z_1(t), \dots, z_m(t)$ in $H_0(f^{-1}(t))$, one can identify each chain $C(t) = \sum_{i=1}^m n_i z_i(t)$ with $n(C) = (n_1, \dots, n_m) \in \mathbb{C}^m$. Similarly, to each vector $v = (v_1, \dots, v_m) \in \mathbb{C}^m$ we associate the chain $C_v := \sum_{i=1}^m v_i z_i(t)$. Then,

$$\int_C g = \langle (g(z_1(t)), \dots, g(z_m(t))), \bar{n}(C(t)) \rangle,$$

where $\langle -, - \rangle$ is the usual scalar product on $\mathbb{C}^m$ and $\bar{n}$ denotes the complex conjugate of n .

Given f and C we search for all g such that $\int_{C(t)} g \equiv 0$. Let G_f be the monodromy group of f and $V(t) \subset H_0(f^{-1}(t))$ the orbit of a chain C of f by G_f . By analytic continuation, the vanishing of $\int_{C(t)} g$ is equivalent to the vanishing of $\int_{\sigma(C(t))} g$, for any $\sigma(C(t)) \in V(t)$. Let $H^0(f^{-1}(t))$ be the dual space to $H_0(f^{-1}(t))$. Hence, from abstract point of view, the tangential center problem is simply the problem of determining $(V(t)^\perp)^*$: the dual space to the orthogonal complement of V . The space $H^0(f^{-1}(t))$ is organized as an m -dimensional $\mathbb{C}[t]$ -module, with multiplication defined by

$$P(t)g(z) = P(f(z))g(z).$$

Let V_r denote r -periodic (mod m) vectors in $\mathbb{Q}^m$ and let $D(f)$ be the set of positive divisors d of $m = \deg(f)$ such that there exists a decomposition $f = \tilde{f} \circ h$, where $\deg(h) = d$. The structure of the G_f -invariant subspaces of $\mathbb{Q}^m$ is determined by Lemma 5.1 of [19].

Lemma 3.1 ([19]). *Any G_f -irreducible invariant subspace of $V(t)$ is of the form*

$$U_r(t) = V_r(t) \cap (V_{r_1}(t)^\perp \cap \dots \cap V_{r_l}(t)^\perp),$$

where $r \in D(f)$ and $\{r_1, \dots, r_l\}$ is a complete set of divisors of m covered by d , that is, they are all the maximal divisors of r in $D(f)$. The subspaces $U_r(t)$ are mutually orthogonal and any G_f -invariant subspace of $\mathbb{C}^m$ is a direct sum of some $U_r(t)$ as above.

For any natural k , w_k denotes the complex vector $(1, \epsilon_m^k, \epsilon_m^{2k}, \dots, \epsilon_m^{(m-1)k}) \in \mathbb{C}^m$, where $\epsilon_m = \exp(2\pi i/m)$ is a primitive m -th root of unity. For $k = 1, \dots, m$ these vectors are orthogonal and form a basis of $\mathbb{C}^m$.

A choice of the chain $C(t)$ determines the corresponding invariant space $V(t)$. Note, moreover, that if a chain $C(t)$ has real coefficients n_j , then for any k the vectors w_k and w_{m-k} simultaneously belong to $V(t)^\mathbb{C}$ or $(V(t)^\perp)^\mathbb{C}$.

Let us recall that $(V_r)^\mathbb{C}$ is generated by the vectors w_k such that m/r divides k . From now on, when necessary we will assume that all $\mathbb{Q}$ -vector spaces $(V(t), V(t)^\perp, V_d, U_d, \text{etc})$ are complexified.

Let $C_k := \bar{w}_k$ and let $\{C_k : k \in S \subset \{1, \dots, m\}\}$ be a basis of the space $V(t)$. Then the solution of the tangential center problem is given by the space generated by the dual basis C_k^* , $k \in \{1, \dots, m\} \setminus S$.

Let P_C denote the characteristic polynomial $P_C(z) = \sum_{j=1}^m n_j z^{j-1}$ and let

$$\Phi_j(z) = \prod_{1 \leq k < j, \gcd(k,j)=1} (z - e^{\frac{2\pi i}{j}k})$$

be the j -th cyclotomic polynomial. Note that $P_C(\epsilon_m^k) = \langle C, \bar{w}_k \rangle = \langle w_k, C \rangle$.

Lemma 3.2. *Let $f(z)$ be a polynomial of degree m and $C(t)$ a chain of f . Then:*

- (1) *The $C[t]$ -module of solutions of the tangential center problem is given by*

$$U^* = U_{d_1}^* \oplus \cdots \oplus U_{d_j}^*,$$

where a basis of U_1^* is given by C_m^* and a basis of U_m^* is given by functions C_k^* , with k relatively prime with m . For any d_j divisor of m , a basis of $U_{d_j}^*$ is given by functions of the form C_k^* , with k a multiple of m/d_j , but not of any other prime factors of d_j , i.e., $\text{g.c.d.}(m, k) = m/d_j$.

- (2) *The subspace U_1^* is a subspace of U^* if and only if $C(t)$ is a cycle. The subspace U_m^* is a subspace of U^* if and only if $C(t)$ is balanced.*
 (3) *The above spaces U_j^* are mutually orthogonal. Their dimensions are given by*

$$\dim_{\mathbb{C}}(U_1^*) = 1, \quad \dim_{\mathbb{C}}(U_m^*) = \phi(m), \quad \dim_{\mathbb{C}}(U_{d_i}^*) = \phi(d_i).$$

- (4) *The set $\{C_j^* : \Phi_{m/\text{g.c.d.}(m,j)}(z) | P_C(z)\}$ is a basis of the space of solutions of the tangential center problem.*

Proof. Assume first that $C(t)$ is a cycle. By definition of a cycle, this means that the chain C_m associated to $w_m = (1, \dots, 1)$ belongs to $V(t)^\perp$. That is, $P_C(\epsilon_m^m) = P_C(1) = 0$. This is equivalent to $\Phi_1(z)$ being a factor of $P_C(z)$ and C_m^* belonging to the space of solutions.

A chain $C(t)$ is unbalanced if for any numeration of the roots corresponding to a permutation cycle at infinity one has $\sum_j n_j \epsilon^j \neq 0$, i.e., the chain C_1 does not belong to $V(t)^\perp$.

Assume now that $C(t)$ is balanced. This means that the chain C_1 also belongs to $V(t)^\perp$, that is, $P_C(\epsilon_m^1) = 0$. By Lemma 3.1, $U_m = V_{r_1}^\perp \cap \cdots \cap V_{r_l}^\perp$, where r_i are all maximal divisors of m in $D(f)$, but strictly smaller than m . This means that m/r_i are all the prime factors of m . In other words, k is coprime with m . Then U_m is generated by C_k , where k is not a multiple of any prime factor of m . Note then that the vector w_1 belongs to U_m , and therefore $U_m \subset V(t)^\perp$. That gives that the functions C_k^* for k coprime with m are in the space of solutions of the center problem and $\Phi_m(z)$ divides $P_C(z)$.

Finally, we calculate the remaining elements of the basis of the solution vector space. They correspond to some irreducible components of $(V^\perp)^*$. By Lemma 3.1 each one of them is of the form $U_d = V_d \cap (V_{r_1}^\perp \cap \cdots \cap V_{r_l}^\perp)$, for some divisor d of m different from 1 and m and a complete set $\{r_1, \dots, r_l\}$ of elements of $D(f)$ covered by d . Similarly as for U_m , now U_d is generated by C_k , where k is a multiple of m/d , but not of other prime factors of d . By duality, for the same k the dual functions C_k^* give a basis of solutions of U_d^* . Besides, that also means that $\Phi_d(z)$ divides $P_C(z)$. $\square$

For some intermediate steps in the solution of the Tangential center problem we shall need to solve not the Tangential center problem but the more general equation

$$\int_{C(t)} g = p(t),$$

where the righthand side is a polynomial $p(t) \in \mathbb{C}[t]$.

Proposition 3.3. *Let $p(t) \in \mathbb{C}[t]$, $p(t) \neq 0$. There exists a solution of $\int_{C(t)} g \equiv p(t)$ if and only if $C(t)$ is not a cycle.*

Moreover, if $C(t)$ is not a cycle, then $\int_{C(t)} g \equiv p(t)$ if and only if $\int_{C(t)} \tilde{g} \equiv 0$, where $\tilde{g} = g - \frac{p \circ f}{\sum n_i}$.

Proof. Assume that $\int_{C(t)} g \equiv p(t)$. Since $p(t)$ is invariant by the action of G_f , then

$$|G_f|p(t) = \sum_{\sigma \in G_f} \sigma(p(t)) = \sum_{\sigma \in G_f} \int_{\sigma(C(t))} g = \frac{|G_f|}{m} \sum_{k=1}^m \left(\sum_{i=1}^m n_i \right) g(z_k(t)).$$

If $C(t)$ is a cycle, it follows $p(t) = 0$, contrary to the assumption.

If $C(t)$ is not a cycle, then $\int_{C(t)} g \equiv p(t)$ is equivalent to

$$0 \equiv \int_{C(t)} g - p(t) = \sum n_i \left(g(z_i(t)) - \frac{p(f(z_i(t)))}{\sum n_i} \right) = \int_{C(t)} \tilde{g}.$$

□

4. SOLUTION FOR 2-TRANSITIVE, MONOMIALS AND CHEBYSHEV POLYNOMIALS

In this section we shall solve the 0-dimensional tangential center problem for the basis of induction, this is, when f is a 2-transitive polynomial, a monomial or a Chebyshev polynomial. We need to consider chains instead of cycles for they appear in the induction process.

First assume that $f(z) = z^m$ or $f(z) = T_m(z)$, where $T_m = \cos(m \arccos(z))$ is the m -th Chebyshev polynomial, with m not necessarily prime. Let $C(t)$ be a balanced cycle of f (with real coefficients). Under the above assumptions we calculate $(V(t)^\perp)^*$ explicitly. The key point of the calculation resides in the fact that dual vectors C_k^* of the chains C_k , or at least the dual space $\text{Vect}(C_k, C_{m-k})^*$ of inseparable spaces $\text{Vect}(C_k, C_{m-k})$, are easily calculated in these cases.

Proposition 4.1. *Let $f(z) = z^m$ (resp. $f(z) = T_m(z)$) be a monomial function (resp. Chebyshev polynomial) and $C(t)$ a balanced chain of f . Then:*

- (1) *The $C[t]$ -module of solutions of the tangential center problem is given by*

$$U^* = U_m^* \oplus U_{d_1}^* \oplus \cdots \oplus U_{d_j}^*,$$

where a basis of U_1^* is given by $g_0(z) = 1$ and a basis of U_m^* is given by the functions $g_k(z) = z^k$ (resp. $g_k(z) = T_k(z)$) for k relatively prime with m . For any d_j divisor of m , a basis of $U_{d_j}^*$ is given by functions of the form $g_k(z) = z^k$ (resp. $g_k(z) = T_k(z)$) for k a multiple of m/d_j , but not of any other prime factors of d_j , i.e., $\text{g.c.d.}(m, k) = m/d_j$.

- (2) *The above spaces U_j^* are mutually orthogonal. Their dimensions are given by*

$$\dim_{\mathbb{C}}(U_1^*) = 1, \quad \dim_{\mathbb{C}}(U_m^*) = \phi(m), \quad \dim_{\mathbb{C}}(U_{d_i}^*) = \phi(d_i).$$

- (3) *The space of solutions of the tangential center problem is generated by $\{g_j(z) = z^j$ (resp. $g_j(z) = T_j(z)\} : \Phi_{m/\text{g.c.d.}(m, j)}(z) | P_C(z)\}$.*

Proof. Let $f(z) = z^m$ or $f(z) = T_m(z)$, $C(t)$ a balanced chain of f and $V(t)$ the orbit of $C(t)$ defined as above. Let $D(m)$ be the set of divisors of m . Note that for any divisor $k \in D(m)$ one has $f = g_{m/k} \circ g_k$, where $g_l(z) = z^l$ (resp. $g_l(z) = T_l(z)$). This shows that in the two particular cases for f a monomial or a Chebyshev polynomial, the set $D(f)$ coincides with $D(m)$, since the decompositions of f are given by the divisors of m . By Lemma 3.2, we now know the complete decomposition of the space of solutions in G_f -invariant spaces, which is the same in both cases. We have to calculate the dual spaces of each of the direct summands of $V(t)^\perp$.

Let $g_l(z) = z^l$ and $z_j(t) = t^{1/m} \epsilon_m^{j-1}$, $j = 1, \dots, m$, for ϵ_m a primitive m -th root of unity. Then

$$\begin{aligned} \int_{C_k} g_l &= \sum_{j=1}^m \overline{w_{k,j}} g_l(z_j(t)) = \sum_{j=1}^m \overline{\epsilon_m^{k(j-1)}} t^{l/m} \epsilon_m^{l(j-1)} \\ &= t^{l/m} \sum_{j=1}^m \overline{\epsilon_m^{k(j-1)}} \epsilon_m^{l(j-1)} = t^{l/m} \langle w_l, w_k \rangle \\ &= m t^{l/m} \delta_{lk}. \end{aligned}$$

This shows that

$$g_l = c_l C_l^*,$$

where c_l is a nonzero constant. The claim in the monomial case follows now from Lemma 3.2.

Consider now the Chebyshev case. Let $T_m(z) = \cos(m \arccos(z))$ be the m -th Chebyshev polynomial, which is a polynomial of degree m . If we take $f(z) = T_m(z)$, then for any $t \in \mathbb{C}$, $f(z) = t$ gives m preimages

$$z_k(t) = (T_m^{-1}(t))_k = \cos\left(\frac{1}{m} \arccos_k(t)\right),$$

where we choose the range of $\arccos_k$ in $[0, 2\pi) + 2(k-1)\pi$ (indeed $\arccos_k(t) = \arccos_1(t) + 2(k-1)\pi$, see Figure 1). Note that a loop of t around infinity transforms $\arccos_k(t)$ into $\arccos_{k+1}(t)$ (see Figure 2).




FIGURE 2. Computation of $z_k(t)$

The Chebyshev polynomials $T_0, \dots, T_{m-1}$ form a basis of the space of polynomials as a $\mathbb{C}[t]$ -module.




FIGURE 3. Monodromic action of a rotation around infinity

If we take $t \in \mathbb{R}$ and denote $\xi(t) = \arccos_1(t) \in \mathbb{R}$, $\epsilon_m = e^{i2\pi/m}$, then for any chain $C(t)$ we obtain

$$\begin{aligned}
 \int_{C(t)} T_j(z) &= \sum_{k=1}^m n_k T_j(z_k) = \sum_{k=1}^m n_k \cos\left(\frac{j}{m} \arccos_k(t)\right) \\
 &= \sum_{k=1}^m n_k \cos\left(\xi(t) \frac{j}{m} + 2(k-1)\pi \frac{j}{m}\right) \\
 &= \sum_{k=1}^m n_k \left(\frac{e^{i\xi(t)\frac{j}{m}}}{2} e^{i2(k-1)\pi \frac{j}{m}} + \frac{e^{-i\xi(t)\frac{j}{m}}}{2} e^{-i2(k-1)\pi \frac{j}{m}} \right) \\
 (4.12) \quad &= \frac{e^{i\xi(t)\frac{j}{m}}}{2} \left(\sum_{k=1}^m n_k \epsilon_m^{j(k-1)} \right) + \frac{e^{-i\xi(t)\frac{j}{m}}}{2} \left(\sum_{k=1}^m n_k \epsilon_m^{-j(k-1)} \right) \\
 &= \frac{e^{i\xi(t)\frac{j}{m}}}{2} P_C(\epsilon_m^j) + \frac{e^{-i\xi(t)\frac{j}{m}}}{2} P_C(\overline{\epsilon_m^j}) \\
 &= \alpha_j(t) P_C(\epsilon_m^j) + \overline{\alpha_j(t)} P_C(\overline{\epsilon_m^j}),
 \end{aligned}$$

where $\alpha_j(t) = \frac{e^{i\xi(t)\frac{j}{m}}}{2}$. If the chain $C(t)$ has real coefficients, this gives

$$\int_{C(t)} T_j(z) = 2\operatorname{Re}(\alpha_j(t) P_C(\epsilon_m^j)),$$

due to the fact that $P_C(z) \in \mathbb{R}[z]$.

Consider in particular the chain C_k . Note that $P_{C_k}(\epsilon_m^j) = \langle w_j, w_k \rangle$. Hence (4.12) gives

$$\int_{C_k(t)} T_j = \alpha_j(t) \langle w_j, w_k \rangle + \overline{\alpha_j(t)} \langle w_{m-j}, w_k \rangle.$$

It follows that $\int_{C_k(t)} T_j = 0$ if $k \notin \{j, m-j\}$.

On the other hand, as our balanced cycle $C(t)$ is real, it follows that the polynomial $P_C(z)$ has real coefficients and hence $\langle w_{m-j}, C \rangle = P_C(\epsilon_m^{m-j}) = P_C(\epsilon_m^{-j}) = \overline{P_C(\epsilon_m^j)} = \overline{P_C(\epsilon_m^j)} = \overline{\langle w_j, C \rangle}$. This means that $w_j \in V(t)^\perp$, if and only if $w_{m-j} \in V(t)^\perp$. It follows that the two dimensional dual space of the space generated by w_j and w_{m-j} is the space generated by T_j and T_{m-j} . The claim now follows as in the monomial case from Lemma 3.2. $\square$

Example 4.2. Let $f(z) = z^6$. Then $D(f) = \{1, 2, 3, 6\}$ and using Lemma 3.1 we get $H_0(f^{-1}(t)) \equiv \mathbb{C}^6 = U_1 \oplus U_2 \oplus U_3 \oplus U_6$, where $U_1 = \langle C_0 \rangle$, $U_2 = \langle C_3 \rangle$, $U_3 = \langle C_2, C_4 \rangle$, $U_6 = \langle C_1, C_5 \rangle$. Let C be a cycle of f , V its orbit and $V^\perp$ the orthogonal complement. By the cycle condition $U_1 \subset V^\perp$.

Assume that the cycle C is balanced. This is equivalent to assuming $U_6 \subset V^\perp$. Now various balanced cycles can be considered. For instance, if C is such that $V = U_2$, then the space of solutions g as a $\mathbb{C}[t]$ -module is generated by $\{1, z, z^2, z^4, z^5\}$. If $V = U_3$, then the solution is generated by $\{1, z, z^3, z^5\}$.

By the Burnside-Schur Theorem, if f is primitive and it is not in the cases above, then it is 2-transitive. The unbalanced case is solved in Theorem 2.3; therefore only the balanced case remains. The following result solves the tangential center problem for a balanced cycle and 2-transitive f .

Proposition 4.3. Let $f \in \mathbb{C}[z]$ be a polynomial with a 2-transitive monodromy group and $C(t) = \sum_{j=1}^m n_j z_j(t)$ a chain of f .

- (1) If $C(t)$ is a balanced chain, then there exists n such that $n_j = n$ for every $1 \leq j \leq m$. In particular if $C(t)$ is a balanced cycle, then it is trivial.
- (2) If $C(t)$ is a balanced chain, then $\int_{C(t)} g \equiv 0$ if and only if

$$\sum_{k=1}^{\deg(g)} s_k g_k = 0,$$

where $g(z) = \sum g_k z^k$ and s_k are given by the Newton-Girard formulae, as

$$s_k = \sum_{i=1}^m z_i^k(t).$$

Proof. (1) Let us assume that $C(t)$ is balanced. It means that $\sum_{j=1}^m n_{\sigma(j)} \epsilon_m^j = 0$, for any $\sigma \in G_f$. Let $H_1 = \{\sigma \in G_f \mid \sigma(1) = 1\}$ be the stabilizer of $z_1(t)$. Then

$$\sum_{\sigma \in H_1} \sum_{j=1}^m n_{\sigma(j)} \epsilon_m^j = 0.$$

That is,

$$(4.13) \quad |H_1| n_1 \epsilon_m + \sum_{j=2}^m \sum_{\sigma \in H_1} n_{\sigma(j)} \epsilon_m^j = 0.$$

Now the assumption that G_f is 2-transitive and H_1 is the stabilizer of $z_1(t)$ implies that H_1 acts transitively on $z_2(t), \dots, z_m(t)$. Hence, for each $j \in \{2, \dots, m\}$ and each $k \in \{2, \dots, m\}$ there is the same number of occurrences of n_k in the sum $\sum_{\sigma \in H_1} n_{\sigma(j)} \epsilon_m^j$. This number is $\frac{|H_1|}{m-1}$ and

$$\sum_{j=2}^m \sum_{\sigma \in H_1} n_{\sigma(j)} \epsilon_m^j = \frac{|H_1|}{m-1} \sum_{k=2}^m n_k \sum_{j=2}^m \epsilon_m^j = \frac{|H_1|}{m-1} \left(-n_1 + \sum_{k=1}^m n_k \right) \sum_{j=2}^m \epsilon_m^j.$$

Observing that $\sum_{j=2}^m \epsilon_m^j = -\epsilon_m$, by Equation (4.13) we get

$$\begin{aligned} 0 &= |H_1| n_1 \epsilon_m - \frac{|H_1|}{m-1} \epsilon_m \left(-n_1 + \sum_{k=1}^m n_k \right) \\ &= |H_1| \epsilon_m \left(n_1 + \frac{n_1}{m-1} - \frac{\sum_{k=1}^m n_k}{m-1} \right) \\ &= \frac{|H_1|}{m-1} \epsilon_m \left(m n_1 - \sum_{k=1}^m n_k \right). \end{aligned}$$

Since $|H_1| \neq 0$ and $\epsilon_m \neq 0$, then

$$n_1 = \frac{\sum_{k=1}^m n_k}{m}.$$

The choice of index 1 was arbitrary. Hence n_j is a constant not depending on j . If $C(t)$ is a cycle, then this constant must be zero.

(2) Let us compute the solutions of $\int_{C(t)} g \equiv 0$, with $g(z) = \sum g_k z^k$, $g_k \in \mathbb{C}$. By (1), we can assume that $C(t) = n \sum z_i(t)$. Then

$$\int_{C(t)} g = n \sum_{i=1}^m g(z_i(t)) = n \sum_{i=1}^m \sum_{k=1}^{\deg(g)} g_k z_i^k(t) = n \sum_{k=1}^{\deg(g)} g_k \sum_{i=1}^m z_i^k(t) \equiv 0,$$

and the solution follows from the Newton-Girard formulae. Recall that Newton-Girard formulae express explicitly $s_k(t) = \sum z_i^k(t)$ in function of the coefficients of the polynomial $f(z) - t$. $\square$

Proof. of Theorem A follows from Propositions 4.1 and 4.3. $\square$

5. MONODROMY GROUP OF IMPRIMITIVE POLYNOMIALS

In the next section we will prove Theorem B and C. We shall assume that Hypothesis 2.8 holds, which, as we prove in this section, will allow us to write the monodromy group of f as a semidirect product defined by the monodromy groups of $\tilde{f}$ and h .

First, we introduce a new numbering in the preimages of t by f . Fix a regular value t of f and take the preimage by $\tilde{f}$ of t . We obtain m/d points, $w_1(t), \dots, w_{m/d}(t)$. For each $w_i(t)$, let $z_{i,j}(t)$, $j = 1, \dots, d$ denote each of the preimages of $w_i(t)$ by h . Then, according to what we saw in Section 1, the blocks of the imprimitivity system associated to $f = \tilde{f} \circ h$ are $B_i = \{z_{i,j}(t) : j = 1, \dots, d\}$, $i = 1, \dots, m/d$. They correspond to rows of circles in Figure 4.

Differentiating $f = \tilde{f} \circ h$, we obtain $f'(z) = \tilde{f}'(h(z))h'(z)$. Thus, critical points of f correspond to either the preimage by h of critical points of $\tilde{f}$ or critical points of h . Let us denote

$$\{z : f'(z) = 0\} = \{a_1, \dots, a_{d(m/d-1)}, b_1, \dots, b_{d-1}\},$$

where $\tilde{f}'(h(a_i)) = 0$, $h'(b_i) = 0$.

Let α_i denote the permutation associated to $f(a_i)$ and β_i the permutation associated to $f(b_i)$. Each permutation α_i (resp. β_i) corresponds to winding counter-clockwise around only one critical value $f(a_i)$ (resp. $f(b_i)$) along a closed path.




FIGURE 4. Regular and critical points/values of imprimitive polynomials

Note that by assumption the paths giving β_i lift to closed paths (loops) based at whatever $w_i(t)$ we take as starting point as they encircle no critical value of $\tilde{f}$. Similar claim is valid for α_i . This gives

$$(5.14) \quad \alpha_k(z_{i,j}) = z_{\alpha_k(i),j}, \quad \beta_k(z_{i,j}) = z_{i,\beta_k(i,j)},$$

so that α_k exchange whole blocks, while β_k only moves elements inside every block.

Thanks to Hypothesis 2.8, we can split the monodromy group G_f of f in terms of the monodromy group of $\tilde{f}$ and a normal subgroup of G_f which contains (a subgroup isomorphic to) the monodromy group of h .

Lemma 5.1. *Assume that Hypothesis 2.8 holds. Let us denote*

$$G_{\tilde{f}} = \langle \alpha_k \rangle, \quad N_h = \langle \alpha_i \beta_k \alpha_i^{-1} : i, k \rangle.$$

Then G_f is the semidirect product $N_h \rtimes_{\phi} G_{\tilde{f}}$ of N_h and $G_{\tilde{f}}$ with respect to $\phi : G_{\tilde{f}} \rightarrow \text{Aut}(N_h)$, $\phi(\alpha) = \phi_{\alpha}$, where $\phi_{\alpha}(\sigma) = \alpha \sigma \alpha^{-1}$.

Proof. First, we define the group semidirect product $N_h \rtimes_{\phi} G_{\tilde{f}}$ as the cartesian product set $N_h \times G_{\tilde{f}}$ with the following operation defined by ϕ :

$$(\sigma, \alpha)(\tilde{\sigma}, \tilde{\alpha}) = (\sigma \phi_{\alpha}(\tilde{\sigma}), \alpha \tilde{\alpha}) = (\sigma \alpha \tilde{\sigma} \alpha^{-1}, \alpha \tilde{\alpha}).$$

Now, since obviously $N_h \cap G_{\tilde{f}} = \{Id\}$, if we prove that any element $\sigma \in G_f$ is written in the form $\sigma = \tau \alpha$, where $\tau \in N_h$ and $\alpha \in G_{\tilde{f}}$, then this decomposition is unique (if $\tau \alpha = \tilde{\tau} \tilde{\alpha}$, then $\tilde{\tau}^{-1} \tau = \tilde{\alpha} \alpha^{-1} \in N_h \cap G_{\tilde{f}} = \{Id\}$ and $\tau = \tilde{\tau}$, $\alpha = \tilde{\alpha}$) and we will have the group isomorphism

$$\begin{aligned} G_f &\longrightarrow N_h \rtimes G_{\tilde{f}} \\ \sigma = \tau \alpha &\mapsto (\tau, \alpha) \end{aligned}$$

Since G_f is generated by the permutations $\{\alpha_i, \beta_j : i = 1, \dots, d(m/d-1), j = 1, \dots, d-1\}$, every element σ in G_f is a finite product of α 's and β 's. In order to write σ as the product of a permutation τ in N_h and a permutation α in $G_{\tilde{f}}$, we group together the α 's and β 's that appears in the writing of σ the following way (we assume without loss of generality that the first group are β 's and the last are α 's):

$$\sigma = \prod_{i \in I_1^{\beta}} \beta_i \prod_{j \in I_2^{\alpha}} \alpha_j \prod_{k \in I_3^{\beta}} \beta_k \dots \prod_{l \in I_r^{\alpha}} \alpha_l,$$

where the set of indices I^{β} are permutations with repetitions of $\{1, \dots, d-1\}$ and the set of indices I^{α} are permutations with repetitions of $\{1, \dots, d(m/d-1)\}$. The first product belongs to N_h , but the second product $\prod_{j \in I_2^{\alpha}} \alpha_j$ is in $G_{\tilde{f}}$, so we multiply the third product $\prod_{k \in I_3^{\beta}} \beta_k$ on the right by $(\prod_{j \in I_2^{\alpha}} \alpha_j)^{-1} \prod_{j \in I_2^{\alpha}} \alpha_j = Id$. Now we rewrite σ as follows

$$\sigma = \prod_{i \in I_1^{\beta}} \beta_i \left(\prod_{j \in I_2^{\alpha}} \alpha_j \prod_{k \in I_3^{\beta}} \beta_k \left(\prod_{j \in I_2^{\alpha}} \alpha_j \right)^{-1} \right) \prod_{j \in I_2^{\alpha}} \alpha_j \dots \prod_{l \in I_r^{\alpha}} \alpha_l,$$

where the first and the second products $\prod_{i \in I_1^{\beta}} \beta_i, \left(\prod_{j \in I_2^{\alpha}} \alpha_j \prod_{k \in I_3^{\beta}} \beta_k \left(\prod_{j \in I_2^{\alpha}} \alpha_j \right)^{-1} \right)$ are in N_h .

We follow this procedure with the new third product until there is no product of β 's left, and in a finite number of times we have σ rewritten as a product of several permutations in N_h and a final one in $G_{\tilde{f}}$, which can be renamed as $\tau \in N_h$ and $\alpha \in G_{\tilde{f}}$. $\square$

Lemma 5.2. *Assume that $f = \tilde{f} \circ h$ satisfies Hypothesis 2.8. Let H_{i_0, j_0} denote the stabilizer of (i_0, j_0) in N_h . Then:*

- (1) $G_{\tilde{f}}$ is isomorphic to the monodromy group of $\tilde{f}$.
- (2) N_h is a normal subgroup of G_f such that for every block B_i of f there exists a subgroup G_i of N_h such that G_i leaves fixed all the elements $(j, k) \notin B_i$. Moreover, G_i is isomorphic to the monodromy group G_h of h .
- (3) The subgroup $\bigcap_{j=1, \dots, d} H_{i_0, j} \subset N_h$ acts transitively on the elements of B_i for every $i \neq i_0$.
- (4) If h is 2-transitive, then H_{i_0, j_0} acts transitively on the other elements of the block B_{i_0} , thus, on $\{(i_0, j) : j \neq j_0\}$.

Proof. (1) and (4) follow easily and (3) is a consequence of (2), since $\bigcap_{j=1, \dots, d} H_{i_0, j}$ contains G_i for every $i \neq i_0$.

To conclude, we prove (2). The group N_h is generated by elements of the form $\tau = \alpha\beta\alpha^{-1}$, with $\alpha \in G_{\tilde{f}}$ and $\beta \in G_h$. By Hypothesis 2.8, β moves elements of at most one block B_i . Since the group $G_{\tilde{f}}$ acts transitively on each column, then for every B_i there exists a subgroup of N_h , which we will call G_i , isomorphic to G_h that moves only the elements of B_i . $\square$

Now, we shall prove that the non-merging hypothesis on critical values implies that for any two decompositions of a polynomial, one of the inner factors factorizes by the other one. We recall that $\tau_\infty = (1, 2, \dots, m) \in G_f$.

Proposition 5.3. *Assume that $f = f_0 \circ h_0$ satisfies Hypothesis 2.8, $f, f_0, h_0 \in \mathbb{C}[z]$. If there exist $f_1, h_1 \in \mathbb{C}[z]$ such that $f = f_1 \circ h_1$, then there exists a polynomial w such that either $h_1 = w \circ h_0$ or $h_0 = w \circ h_1$.*

Proof. Let B_0 be a block associated to the decomposition $f = f_0 \circ h_0$ such that $1 \in B_0$, and let B_1 a block associated to $f = f_1 \circ h_1$ such that $1 \in B_1$. Then either $B_1 \subset B_0$ or there exists $i \in B_1 \setminus B_0$.

If $B_1 \subset B_0$, then there exists $w \in \mathbb{C}[z]$ such that $w(h_1) = h_0$: we know that $B_0 = \{j \in \{1, \dots, m\} : h_0(z_j) = h_0(z_j)\}$ and $B_1 = \{k \in \{1, \dots, m\} : h_1(z_1) = h_1(z_k)\}$. Their stabilizers $G_{B_0} = \{\sigma \in G_f : \sigma(B_0) = B_0\}$ and $G_{B_1} = \{\sigma \in G_f : \sigma(B_1) = B_1\}$ verify that $G_{B_1} \subset G_{B_0}$ and, therefore, $L^{G_{B_0}} = \mathbb{C}(h_0(z_1)) \subset L^{G_{B_1}} = \mathbb{C}(h_1(z_1))$, where $L = \mathbb{C}(z_1(t), \dots, z_m(t))$. As a consequence, there exists a rational function w , which we can assume polynomial, such that $h_0 = w \circ h_1$.

If there exists $i \in B_1 \setminus B_0$, then $\tau_\infty^{i-1}(B_0) \subset B_1$, and there exists $w \in \mathbb{C}[z]$ such that $w(h_0) = h_1$: since $\tau_\infty^{i-1}(1) = i \in \tau_\infty^{i-1}(B_0)$, this one is a block of the imprimitivity system associated to the decomposition $f = f_0 \circ h_0$ different from B_0 , then $1 \notin \tau_\infty^{i-1}(B_0)$. By Lemma 5.2 (3), we can assure that the stabilizer H_1 (in G_f) of 1 is transitive on every other block different from B_0 . Therefore, H_1 is transitive on $\tau_\infty^{i-1}(B_0)$ and the orbit of i by H_1 contains $\tau_\infty^{i-1}(B_0)$. Since every permutation in H_1 leaves B_1 fixed and $i \in B_1$, $\tau_\infty^{i-1}(B_0) \subset B_1$ and following a similar argument as in the previous case, there exist $w \in \mathbb{C}[z]$ such that $h_1 = w \circ h_0$. $\square$

Proposition 5.3 implies that when we have chains of decompositions satisfying Hypothesis 2.8, then we have uniqueness.

Proof of Proposition 2.9. We shall prove it by induction on the number $\tilde{d}$ of factors in the decomposition of f .

First, assume that $\tilde{d} = 0$ and $f = \tilde{f}_0$ is 2-transitive, Chebyshev or a monomial, and let $f = f_0 \circ \dots \circ f_d$ be a decomposition of f for some polynomials $f_0, \dots, f_d$ such that every f_k is either 2-transitive, Chebyshev or a monomial, and that Hypothesis 2.8 holds for every $f_0 \circ \dots \circ f_k = (f_0 \circ \dots \circ f_{k-1}) \circ f_k$. Since 2-transitive polynomials are primitive and monomials or Chebyshev polynomials can not decompose satisfying Hypothesis 2.8, then $d = 0$.

Now, assume that $f = f_0 \circ \dots \circ f_d$ for some polynomials $f_0, \dots, f_d$ such that every f_k is either 2-transitive, Chebyshev or a monomial, and that Hypothesis 2.8 holds for every $f_0 \circ \dots \circ f_k = (f_0 \circ \dots \circ f_{k-1}) \circ f_k$.

If $f = \tilde{f}_0 \circ \dots \circ \tilde{f}_{\tilde{d}}$ is another decomposition of f for some polynomials $\tilde{f}_0, \dots, \tilde{f}_{\tilde{d}}$ such that every $\tilde{f}_k$ is either 2-transitive, Chebyshev or a monomial, by Proposition 5.3 there exists $w \in \mathbb{C}[z]$ such that either $f_d = w \circ \tilde{f}_{\tilde{d}}$ or $\tilde{f}_{\tilde{d}} = w \circ f_d$.

Assume that $\tilde{f}_{\tilde{d}} = w \circ f_d$ for some non-linear w . In particular this implies that w, f_d are both monomials or both Chebyshev polynomials. Then

$$f_0 \circ \dots \circ f_{d-1} = \tilde{f}_0 \circ \dots \circ \tilde{f}_{\tilde{d}-1} \circ w.$$

Thus, Hypothesis 2.8 does not hold, since $\tilde{f}_0 \circ \dots \circ \tilde{f}_{\tilde{d}-1} \circ w$ and f_d share a critical value.

Therefore, we may assume that $f_d = w \circ \tilde{f}_{\tilde{d}}$. But in this case

$$f_0 \circ \dots \circ f_{d-1} \circ w = \tilde{f}_0 \circ \dots \circ \tilde{f}_{\tilde{d}-1}.$$

Note that $f_0 \circ \dots \circ f_{d-1} \circ w$ still satisfies Hypothesis 2.8 since the critical values of w are a subset of those of f_d . We conclude by induction.

If Hypothesis 2.8 holds for every $\tilde{f}_0 \circ \dots \circ \tilde{f}_k = (\tilde{f}_0 \circ \dots \circ \tilde{f}_{k-1}) \circ \tilde{f}_k$, interchanging $\{f_k\}$ and $\{\tilde{f}_k\}$ in the previous arguments we conclude. $\square$

A decomposition of f in primitive polynomials is not unique in general. To get uniqueness, we have to regroup successive monomials or Chebyshev factors (or factors linearly equivalent to one of them) that commute (see [18]). Ritt's Theorem states that if a polynomial f admits two different decompositions, then there exists monomials or a Chebyshev factors (or factor linearly equivalent to one of the previous types) in a decomposition of f into prime polynomials that commute. Note for instance that $z^6 = (z^2)^3 = (z^3)^2$. In this section we have studied the converse problem, thus, given a decomposition of $f = \tilde{f} \circ h$, Hypothesis 2.8 ensures that any other decomposition of f is a further decomposition of $\tilde{f}$ and h , or, in other words, there exists no commuting factors between $\tilde{f}$ and h .

6. PROOF OF THEOREMS B AND C

Let $C(t)$ be a balanced cycle of an imprimitive polynomial $f = \tilde{f} \circ h$, with h 2-transitive, Chebyshev or monomial. According to the type of h , we deduce first some information on how the cycle $C(t)$ is positioned with respect to the imprimitivity system defined by h .

Proposition 6.1. *Assume that $f = \tilde{f} \circ h$ satisfies Hypothesis 2.8. Let $C(t)$ be a balanced cycle of f and let $\mathcal{B}_h = \{B_1, \dots, B_{m/d}\}$ denote the imprimitivity system corresponding to h .*

- (1) *If h is 2-transitive, then n_k is constant for $k \in B_j$, for every $B_j \in \mathcal{B}_h$.*
- (2) *If $h(z) = z^d$ or $h(z) = T_d(z)$, then the restriction of $C(t)$ to each block of $\mathcal{B}_h$ is balanced.*

Proof. (1) Let us fix i_0, j_0 and let us denote $H_0 = H_{i_0, j_0} \subset G_f$ the stabilizer of z_{i_0, j_0} . For every preimage $z_{i, j}(t)$ of t by f , let us denote by $k(i, j)$ its position when we enumerate them so that $\tau_\infty = (z_1(t), z_2(t), \dots, z_m(t))$. Besides, $z_{i, j}(t) = z_{k(i, j)}(t)$.

Therefore, since $C(t)$ is balanced,

$$\begin{aligned} 0 &= \sum_{\tau \in H_0} \sum_{i,j} n_{\tau(i,j)} \epsilon_m^{k(i,j)} \\ &= |H_0| n_{i_0, j_0} \epsilon_m^{k(i_0, j_0)} + |H_0| \left(\frac{\sum_{j=1, j \neq j_0}^d n_{i_0, j}}{d-1} \sum_{j=1, j \neq j_0}^d \epsilon_m^{k(i_0, j)} \right) + \\ &\quad |H_0| \left(\sum_{i=1, i \neq i_0}^{m/d} \frac{\sum_{j=1}^d n_{i, j}}{d} \sum_{j=1}^d \epsilon_m^{k(i, j)} \right), \end{aligned}$$

where the second summand is due to Lemma 5.2 (4), since h is 2-transitive, and the last summand is due to Lemma 5.2 (3). Now, observe that $\sum_{j=1}^d \epsilon_m^{k(i, j)} = 0$ for every i , since it is the sum of all the powers of ϵ_m corresponding to a block B_i , whose elements are the residue class mod m/d for some $l \in \{1, \dots, d\}$ (see, for instance, [19, Lemma 3.1]), that is, $\sum_{j=1}^d \epsilon_m^{k(i, j)} = \sum_{j=1}^d \epsilon_m^{l+jm/d} = \epsilon_m^l \sum_{j=1}^d \epsilon_m^{jm/d} = \epsilon_m^l \sum_{j=1}^d \epsilon_d^j = 0$. Therefore,

$$\begin{aligned} 0 &= |H_0| \left(n_{i_0, j_0} \epsilon_m^{k(i_0, j_0)} + \frac{\sum_{j=1, j \neq j_0}^d n_{i_0, j}}{d-1} (-\epsilon_m^{k(i_0, j_0)}) \right) \\ &= |H_0| \epsilon_m^{k(i_0, j_0)} \left(n_{i_0, j_0} (1 + 1/(d-1)) - \frac{\sum_{j=1}^d n_{i_0, j}}{d-1} \right). \end{aligned}$$

Thus, n_{i_0, j_0} does not depend on j_0 . Therefore $n_{i_0, j}$ is constant in the block B_{i_0} .

(2) Assume now that $h(z) = z^d$, or $h(z) = T_d(z)$. Let $H_0 = \bigcap_{j=1, \dots, d} H_{i_0, j}$. The elements $\{(i, j)\}_{j=1, \dots, \deg(\tilde{f})}$ with $i \neq i_0$ fixed are moved transitively by Lemma 5.2 (3). Therefore, using similar arguments as in (1),

$$0 = \sum_{\tau \in H_0} \sum_{i, j} n_{\tau(i, j)} \epsilon_m^{k(i, j)} = |H_0| \sum_{j=1}^d n_{i_0, j} \epsilon_m^{k(i_0, j)}.$$

Then, $C(t)$ restricted to the block B_{i_0} (the i_0 -th h -invariant part of $C(t)$) is balanced. $\square$

Proof of Theorem B. By Proposition 6.1, $C(t)$ is constant along the blocks of the imprimitivity system corresponding to h , $\mathcal{B}_h = \{B_1, \dots, B_{m/d}\}$. Therefore,

$$\int_{C(t)} g = \sum_i n_i g(z_i(t)) = \sum_{i=1}^{m/d} n_i \sum_{k \in B_i} g(z_k(t)).$$

Let us observe that $\sum_{k \in B_i} g(z_k(t))$ is invariant by $G_{B_i} = \{\sigma \in G_f : \sigma(B_i) = B_i\}$. Then, if we denote $L = \mathbb{C}(z_1(t), \dots, z_m(t))$, $\sum_{k \in B_i} g(z_k(t)) \in L^{G_{B_i}} = \mathbb{C}(h(z_k(t)))$ for every $k \in B_i$. Let us denote $w_i(t) = h(z_k(t))$, which is constant in $k \in B_i$; then $w_i(t)$ is one of the m/d preimages of t by $\tilde{f}$. Therefore, $\sum_{k \in B_i} g(z_k(t)) = \tilde{g}_i(w_i(t))$ for some rational function $\tilde{g}_i(z)$, which we can assume is a polynomial. Since every subgroup G_{B_i} of G_f is conjugated to another G_{B_j} , it can easily be proved that $\tilde{g}_i$ does not depend on the block B_i , that is, there is only one polynomial $\tilde{g}(z)$ such that

$$(6.15) \quad \sum_{k \in B_i} g(z_k(t)) = \tilde{g}(w_i(t)), \quad i = 1, \dots, m/d.$$

Then, since $h(C(t)) = \sum_{i=1}^{m/d} dn_i w_i(t)$,

$$\int_{C(t)} g = \sum_{i=1}^{m/d} n_i \tilde{g}(w_i(t)) = \frac{\int_{h(C(t))} \tilde{g}}{d}.$$

Since $z_k(t)$ for $k \in B_i$ are the preimages of w_i by h , Equation (6.15) can be rewritten as

$$\sum_{k \in B} g(z_k(w)) = \tilde{g}(w).$$

Replacing g by its linear expansion in terms of the basis $\{1, z, \dots, z^{d-1}, h(z), zh(z), \dots, z^{d-1}h(z), h^2(z), \dots\}$ of $\mathbb{C}[z]$, namely $g(z) = \sum_{i=0}^{d-1} z^i g_i(h(z))$, we get

$$\begin{aligned} \tilde{g}(w) &= \sum_{k \in B} \sum_{i=0}^{d-1} z_k^i(w) g_i(h(z_k(w))) = \sum_{k \in B} \sum_{i=0}^{d-1} z_k^i(w) g_i(w) \\ &= \sum_{i=0}^{d-1} \left(\sum_{k \in B} z_k^i(w) \right) g_i(w) = \sum_{i=0}^{d-1} s_i g_i(w), \end{aligned}$$

where s_i can be obtained by the Newton-Girard formulae applied to h . $\square$

Proof of Theorem C. Assume that $f = \tilde{f} \circ h$, $h(z) = z^d$ or $h(z) = T_d(z)$ and Hypothesis 2.8 is verified. Let $C(t)$ be a balanced cycle of f . Assume that

$$\int_{C(t)} g \equiv 0,$$

that is, $\sum_{i,j} n_{i,j} g(z_{i,j}(t)) = 0$. Fix $i_0 \in \{1, \dots, m/d\}$. Let H_{i_0} be the stabilizer of (i_0, j) , $j = 1, \dots, d$ in N_h . By Lemma 5.2 (3), the elements (i, j) , $j = 1, \dots, d$ with $i \neq i_0$ are moved transitively by H_{i_0} . Therefore,

$$\begin{aligned} 0 &= \sum_{\tau \in H_{i_0}} \sum_{i,j} n_{\tau(i,j)} g(z_{i,j}(t)) \\ &= |H_{i_0}| \sum_{j=1}^d n_{i_0,j} g(z_{i_0,j}(t)) + \frac{|H_{i_0}|}{d-1} \sum_{i=1, i \neq i_0}^{m/d} \left(\sum_{j=1}^d n_{i,j} \right) \left(\sum_{j=1}^d g(z_{i,j}(t)) \right). \end{aligned}$$

Let us observe that $\sum_{j=1}^d g(z_{i,j}(t))$ is invariant by $G_{B_i} = \{\sigma \in G_f : \sigma(B_i) = B_i\}$, for $i \neq i_0$. As a consequence, if we denote $L = \mathbb{C}(z_1(t), \dots, z_m(t))$, we obtain that $\sum_{j=1}^d g(z_{i,j}(t)) \in L^{G_{B_i}} = \mathbb{C}(h(z_{i,j}(t))) = \mathbb{C}(w_i(t))$, where $w_i(t) = h(z_{i,j}(t))$ for $j = 1, \dots, d$. By Lüroth Theorem, there exists a polynomial $\tilde{g}(w)$ (which as in previous proof does not depend on i) such that

$$\sum_{j=1}^d g(z_{i,j}(t)) = \tilde{g}(w_i(t))$$

and, consequently,

$$\sum_{j=1}^d n_{i_0,j} g(z_{i_0,j}(t)) = \frac{-1}{d-1} \sum_{i=1, i \neq i_0}^{m/d} \left(\sum_{j=1}^d n_{i,j} \right) \tilde{g}(w_i(t)).$$

Replacing in the equation $\int_{C(t)} g = 0$,

$$\begin{aligned} 0 &= \sum_{i=1}^{m/d} \sum_{j=1}^d n_{i,j} g(z_{i,j}(t)) = \frac{-1}{d-1} \sum_{i_0=1}^{m/d} \sum_{i=1, i \neq i_0}^{m/d} \left(\sum_{j=1}^d n_{i,j} \right) \tilde{g}(w_i(t)) \\ &= -\frac{\frac{m}{d}-1}{d-1} \sum_{i=1}^{m/d} \left(\sum_{j=1}^d n_{i,j} \right) \tilde{g}(w_i(t)). \end{aligned}$$

Thus,

$$(6.16) \quad \int_{h(C(t))} \tilde{g} = 0.$$

On the other hand,

$$\begin{aligned} \sum_{j=1}^d n_{i_0,j} g(z_{i_0,j}(t)) &= \frac{-1}{d-1} \sum_{i=1, i \neq i_0}^{m/d} \left(\sum_{j=1}^d n_{i,j} \right) \tilde{g}(w_i(t)) \\ &= \frac{1}{d-1} \left(\sum_{j=1}^d n_{i_0,j} \right) \tilde{g}(w_{i_0}(t)). \end{aligned}$$

In consequence, putting

$$(6.17) \quad p_{i_0}(w) := \frac{\sum_{j \in B_{i_0}} n_j}{d-1} \tilde{g}(w) \in \mathbb{C}[w]$$

for every i_0 , g is a solution of

$$(6.18) \quad \int_{\tilde{C}_{i_0}(w)} g = p_{i_0}(w), \quad 1 \leq i_0 \leq m/d.$$

Recall that $\tilde{C}_k(w)$ is a chain of h and the right-hand part of (6.18) is a polynomial. Then, by Proposition 3.3 a particular solution of (6.18) is given by

$$\frac{(p_{i_0} \circ h)(z)}{\sum_{i \in B_{i_0}} n_i} = \frac{\tilde{g}(h(z))}{d-1},$$

and the general solution is given by adding general solutions of $\int_{\tilde{C}_{i_0}(w)} u = 0$. $\square$

REFERENCES

- [1] A. Álvarez, J.L. Bravo, C. Christopher, *Some results on the trigonometric moment problem*, preprint arXiv:1108.0172v2 20 Sept. 2011
- [2] A. Álvarez, J.L. Bravo, P. Mardešić, *Vanishing Abelian integrals on zero-dimensional cycles*, arXiv: 1101.1777v2 [math.DS] 17 Jun. 2011 ((v1)10 Jan. 2011).
- [3] P. Bonnet, A. Dimca, *Relative differential forms and complex polynomials*, Bull. Sci. math., 124-7 (2000) 557–571.
- [4] M. Briskin, J.P. Françoise, Y. Yomdin, *The Bautin ideal of the Abel equation* Nonlinearity **11** (1998) 41–53.
- [5] M. Briskin, J.P. Françoise, Y. Yomdin, Y., *Center conditions, compositions of polynomials and moments on algebraic curve*, Ergod. Theor. Dyn. Syst. **19**(5) (1999) 1201–1220.
- [6] M. Briskin, J.P. Françoise, Y. Yomdin, *Center conditions II: Parametric and model center problems*, Isr. J. Math. **118** (2000) 61–82.
- [7] M. Briskin, J.P. Françoise, Y. Yomdin, *Center conditions III: Parametric and model center problems*, Israel J. Math. **118** (2000) 83–108.
- [8] M. Briskin, N. Roytvarf, Y. Yomdin, *Center conditions at infinity for Abel differential equation*, Annals of Math., 172 (2010) 437–483
- [9] A. Cima, A. Gasull, F. Mañosas, *Strongly persistent centers for trigonometric Abel equations*, to appear in Qual. Theory Dyn. Syst.

- [10] C. Christopher, P. Mardešić, *The monodromy problem and the tangential center problem*, Funkts. Anal. Prilozh., 44-1 (2010) 27–43.
- [11] J.D. Dixon, B. Mortimer, *Permutation groups*, Graduate Texts in Mathematics **163**, Springer-Verlag, New-York (1996).
- [12] H. Dulac, *Détermination et intégration d'une certaine classe d'équations différentielle ayant pour point singulier un centre*, Bull. Sci. Math. Sér. (2) **32** (1) (1908), 230–252.
- [13] L. Gavrilov, *Petrov modules and zeros of Abelian integrals*, Bull. Sci. Math., 122-8 (1998) 571–584.
- [14] L. Gavrilov, F. Pakovich, *Moments on Riemann surfaces and hyperelliptic Abelian integrals*, preprint arXiv:1107.3029v1 15 Jul. 2011.
- [15] Yu. Ilyashenko, *Appearance of limit cycles by perturbation of the equation $dw/dz = -R_z/R_w$, where $R(z; w)$ is a polynomial*, Mat. Sbornik (New Series) 78-3 (1969) 360–373.
- [16] W. Kapteyn, *New investigations on the midpoints of integrals of differential equations of the first degree*, Nederl. Akad. Wetensch. Verslag Afd. Natuurk. **20** (1912), 1354-1365; **21**, 27-33. (Dutch)
- [17] P. Müller, M.E. Zieve, *On Ritt's polynomials decomposition theorems*, preprint.
- [18] F. Pakovich, *Generalized “second Ritt theorem” and explicit solutions of the polynomial moment problem*, preprint.
- [19] F. Pakovich, M. Muzychuk, *Solution of the polynomial moment problem*, Proc. Lond. Math. Soc., 99-3 (2009) 633–657.
- [20] J.F. Ritt, *Prime and composite polynomials*, Trans. Amer. Math. Soc., 23 (1922) 51–66.